

Erratum to: “Banach Spaces without Local Unconditional Structure”

Ryszard A. Komorowski Nicole Tomczak-Jaegermann

7 July, 1996

Abstract

This note contains a corrected proof of the main result (which remains unchanged) from [K-T]. It was recently observed that an argument in a basic technical criterium has a gap.

The first named author recently observed that the paper by Komorowski and Tomczak [K-T], contains a gap. Here we shall correct the formulation of a general result (Theorem 2.1 in [K-T]) and then we shall complete the proof of the main application to the homogeneous Banach space problem (Theorem 3.1 in [K-T]), that is now necessary in view of modifications in the aforementioned general statement. Corollaries 4.3, 4.4 and 4.5 do not require any changes.

This note is not self-contained. However, in order to make it possible for the readers a little familiar with [K-T], to read it without constantly referring to the original paper, we decided not to cut the arguments to bare minimum, and occasionally give some additional background information.

The gap is in the proof of the fundamental criterium from [K-T] (Proposition 1.1). One deals there with a Banach space Y which has a 2-dimensional Schauder decomposition $\{Z_k\}_k$. Assuming that Y has cotype $q < \infty$ and l.u.st., one constructs a block diagonal operator T , which was claimed to be bounded on subspaces Y_K for which the decomposition $\{Z_k\}_{k \in K}$ was unconditional. In fact, we can estimate this norm only if the subspace is complemented by a natural projection. To be more precise, for a subset $K \subset \mathbb{N}$ we define $Q_K : Y \rightarrow Y_K$ by $Q_K(\sum_k z_k) = \sum_{k \in K} z_k$. Then the correct formulation

of Proposition 1.1 states, in condition (ii), that if the decomposition $\{Z_k\}_{k \in K}$ of Y_K is C -unconditional and $\|Q_K\| \leq C'$, then

$$\|T|_{Y_K}: Y_K \rightarrow Y_K\| \leq C^2 C' \psi \text{ l.u.st.}(Y), \quad (1)$$

where ψ is the same function as in [K-T].

The proof uses the same operator T as in [K-T], and the estimate is completely clear from the original proof, once we notice that letting $P_k: Y \rightarrow Z_k$ to denote the natural projection onto Z_k , for $k = 1, 2, \dots$, we trivially have $P_k = P_k Q_K$ for every $k \in K$, and thus, for arbitrary signs $\varepsilon_k = \pm 1$ we have

$$\left\| \sum_{k \in K} \varepsilon_k P_k : Y \rightarrow Y_K \right\| = \left\| \sum_{k \in K} \varepsilon_k P_k Q_K \right\| \leq C C'.$$

Remark Note that if the decomposition $\{Z_k\}_k$ of Y is unconditional, the natural projections Q_K are automatically bounded for all $K \subset \mathbb{N}$. In this case the proof of Proposition 1.1 in [K-T] was correct, so Corollaries 4.3, 4.4 and 4.5 do not require any changes.

The remaining part of this note heavily depends on technical notation from the end of Section 1 of [K-T]. For the reader convenience we copy it here.

If $\Delta = \{A_m\}_m$ and $\Delta' = \{A'_m\}_m$ are partitions of $\mathbb{N}$, we say that $\Delta \succ \Delta'$, if there exists a partition $\mathcal{J}(\Delta', \Delta) = \{J_m\}_m$ of $\mathbb{N}$ such that

$$\min J_m < \min J_{m+1} \quad \text{and} \quad A'_m = \bigcup_{j \in J_m} A_j \quad \text{for } m = 1, 2, \dots \quad (2)$$

In such a situation, for $m = 1, 2, \dots$, $\mathcal{K}(A'_m, \Delta)$ denotes the family

$$\mathcal{K}(A'_m, \Delta) = \{K \subset A'_m \mid |K \cap A_j| = 1 \text{ for } j \in J_m\}. \quad (3)$$

Finally, if $\Delta_i = \{A_{i,m}\}_m$, for $i = 1, 2, \dots$, is a sequence of partitions of $\mathbb{N}$, with $\Delta_1 \succ \dots \succ \Delta_i \succ \dots$, we set, for $m = 1, 2, \dots$ and $i = 2, 3, \dots$

$$\mathcal{K}_{i,m} = \mathcal{K}(A_{i,m}, \Delta_{i-1}). \quad (4)$$

Now Theorem 2.1 from [K-T] requires an obvious extra assumption of boundness of natural projections. Although the statement of this theorem is rather lengthy, we recall it here (in a corrected form) for sake of future references.

Theorem ★2.1: Let $X = F_1 \oplus \dots \oplus F_4$ be a direct sum of Banach spaces of cotype r , for some $r < \infty$, and let $\{f_{i,l}\}_l$ be a normalized monotone Schauder basis in F_i , for $i = 1, \dots, 4$. Let $\Delta_1 \succ \dots \succ \Delta_4$ be partitions of $\mathbb{N}$, $\Delta_i = \{A_{i,m}\}_m$ for $i = 1, \dots, 4$. Assume that there exist $C, C' \geq 1$ such that for every $K \in \mathcal{K}_{i,m}$ with $i = 2, 3, 4$ and $m = 1, 2, \dots$, the basis $\{f_{s,l}\}_{l \in K}$ in $F_s|_K$ is C -unconditional, and the natural projection $R_K^{(s)}: F_s \rightarrow F_s|_K$ has the norm $\|R_K^{(s)}\| \leq C'$, for $s = 1, \dots, 4$; moreover, there is $\tilde{C} \geq 1$ such that for $i = 1, 2, 3$ and $m = 1, 2, \dots$ we have

$$\|I: F_i|_{A_{i,m}} \rightarrow F_{i+1}|_{A_{i,m}}\| \leq \tilde{C}. \quad (5)$$

Assume finally that one of the following conditions is satisfied:

- (i) there is a sequence $0 < \delta_m < 1$ with $\delta_m \downarrow 0$ such that for every $i = 1, 2, 3$ and $m = 1, 2, \dots$ and every $K \in \mathcal{K}_{i+1,m}$ we have

$$\|I: D(F_1 \oplus \dots \oplus F_i)|_K \rightarrow F_{i+1}|_K\| \geq \delta_m^{-1}; \quad (6)$$

- (ii) there is a sequence $0 < \delta_m < 1$ with $\delta_m \downarrow 0$ and $\sum_m \delta_m^{1/2} = \gamma < \infty$ such that for every $i = 1, 2, 3$ and $m = 1, 2, \dots$ and every $K \in \mathcal{K}_{i+1,m}$ we have

$$\|I: F_{i+1}|_K \rightarrow F_i|_K\| \geq \delta_m^{-1}. \quad (7)$$

Then there exists a subspace Y of X without local unconditional structure, but which still admits a Schauder basis.

Proof The only addition to the original proof is to check that the assumption for $\|Q_K\|$ required in the fundamental criterium is now satisfied. Since spaces Z_k are spanned by vectors of the form

$$\begin{aligned} x_k &= \alpha_{1,k} f_{1,k} + \dots + \alpha_{4,k} f_{4,k}, \\ y_k &= \alpha'_{1,k} f_{1,k} + \dots + \alpha'_{4,k} f_{4,k}, \end{aligned}$$

then we have $Q_K(\sum_k z_k) = \sum_{s=1}^4 R_K^{(s)}\left(\sum_k (t_k \alpha_{s,k} + t'_k \alpha'_{s,k}) f_{s,k}\right)$, for $z = \sum_k z_k = \sum_k (t_k x_k + t'_k y_k)$, thus $\|Q_K\| \leq 4 \max_s \|R_K^{(s)}\|$. $\square$

The remaining part of this note contains the proof of Theorem 3.1 from [K-T], which says:

Theorem 3.1: *Let X be a Banach space with an unconditional basis and of cotype r , for some $r < \infty$. If X does not contain a subspace isomorphic to l_2 then there exists a subspace Y of X without local unconditional structure, which admits a Schauder basis.*

Let $\{e_l\}_l$ be a 1-unconditional basis in X . Passing to a subsequence we may assume without loss of generality that $\{e_l\}_l$ generates a spreading model $\{u_l\}_l$. We distinguish two main cases.

- (I) either no sequence of disjointly supported blocks of $\{e_l\}_l$ of length ≤ 3 with constant coefficients is equivalent to $\{u_l\}_l$,
- (II) or the basis $\{e_l\}_l$ is 1-subsymmetric.

If Case I does not hold, then there exist disjointly supported blocks $\{w_l\}_l$ equivalent to $\{u_l\}_l$; so by renorming we get into Case II for $X_1 = \overline{\text{span}} \{w_l\}_l$. Case I is new, compared with [K-T]. Case II has the same proof as in [K-T]. The additional information that the basis is subsymmetric implies that equally distributed vectors spanning ℓ_2^n 's are given by constant coefficient blocks (see Lemma 3.2 in [K-T] and the remark after). Then the required complementation follows from the classical fact that in a subsymmetric space subspaces spanned by constant coefficient consecutive blocks are well complemented.

Case I is the content of the following proposition proved by B. Maurey and N. Tomczak in May 1993 (unpublished).

Proposition I: *Let X be a Banach space of cotype r , for some $r < \infty$. Assume that X has a 1-unconditional basis $\{e_l\}_l$ which generates a spreading model $\{u_l\}_l$. If no disjointly supported blocks of $\{e_l\}_l$ of length ≤ 3 with constant coefficients are equivalent to $\{u_l\}_l$, then there exists a subspace Y of X without local unconditional structure, which admits a 2-dimensional unconditional decomposition*

Proof Denote the span of $\{u_l\}_l$ by E . First observe, by a compactness argument, that if no subsequence of $\{e_l\}_l$ dominates (resp. is dominated by) $\{u_l\}_l$ and if $\{A_m\}_m$ is a partition of $\mathbb{N}$ into finite sets then for every D there exists M such that whenever K is a subset of $\mathbb{N}$ such that $|K \cap A_m| = 1$ for $1 \leq m \leq M$ then $\|I : X|_K \rightarrow E|_K\| \geq D$ (resp. $\|I : E|_K \rightarrow X|_K\| \geq D$).

By passing to subsequences of $\{e_l\}_l$ we may assume that

- (Ia) either $\|I: E|_L \rightarrow X|_L\| = \infty$ for all infinite subsequences L of N ,
 (Ib) or $\|I: E \rightarrow X\| < \infty$.

Partition the whole basis $\{e_l\}_l$ into four infinite sets $\{e_{i,l}\}_l$ for $i = 1, \dots, 4$. Of course E is the spreading model for each $\{e_{i,l}\}_l$ ($i = 1, \dots, 4$). In the proof we shall use a shorthand notation $\sim$ to denote that two sequences are 2-equivalent.

Case (Ia): Set $f_{1,l} = e_{1,l}$ for all l and let $\Delta_1 = \{A_{1,m}\}_m$ consists of singletons. Let $F_1 = \overline{\text{span}}[f_{1,l}]_l$. Let $\Delta_2 = \{A_{2,m}\}_m$ be a partition of N into successive intervals such that $\|I: E|_{A_{2,m}} \rightarrow F_1|_{A_{2,m}}\| \geq 2^{2m}$ for all m . Define $f_{2,l}$ for $l \in A_{2,m}$ by induction in m . For $l \in A_{2,1}$ set $f_{2,l} = e_{2,s_1+l}$, where s_1 is so large that $\{f_{2,l}\}_{l \in A_{2,1}} \sim \{u_l\}_{l=1}^{|A_{2,1}|}$. Then having defined $f_{2,l}$ for all $l \in A_{2,m}$ set, for $l \in A_{2,m+1}$, $f_{2,l} = e_{2,s_{m+1}+l}$, where $s_{m+1} > s_m + |A_{2,m}|$ is so large that $\{f_{2,l}\}_{l \in A_{2,m+1}} \sim \{u_l\}_{l=1}^{|A_{2,m+1}|}$. In particular, $\{f_{2,l}\}_l$ is a subsequence of $\{e_{2,l}\}_l$. Let $F_2 = \overline{\text{span}}[f_{2,l}]_l$.

By our initial remark we can define by induction a sequence $0 = M_0 < M_1 < \dots < M_j < \dots$ such that for every $j = 1, 2, \dots$ whenever $K \subset N$ satisfies $|K \cap A_{2,m}| = 1$ for $M_{j-1} < m \leq M_j$, then $\|I: E|_K \rightarrow F_2|_K\| \geq 2^{2j}$. Then set $A_{3,j} = \bigcup_{m=M_{j-1}+1}^{M_j} A_{2,m}$ for $j = 1, 2, \dots$. Define $\{f_{3,l}\}_l$ as a subsequence of $\{e_{3,l}\}_l$ such that $\{f_{3,l}\}_{l \in A_{3,j}} \sim \{u_l\}_{l=1}^{|A_{3,j}|}$, and set $F_3 = \overline{\text{span}}[f_{3,l}]_l$.

Finally, using the same construction once more for the partition $\Delta_3 = \{A_{3,j}\}_j$, find a partition $\Delta_4 = \{A_{4,n}\}_n$ with $\Delta_4 \prec \Delta_3$ such that for every $K \in \mathcal{K}_{4,n}$ we have $\|I: E|_K \rightarrow F_3|_K\| \geq 2^{2n}$ (for $n = 1, 2, \dots$). Then define $\{f_{4,l}\}_l$ as a subsequence of $\{e_{4,l}\}_l$ such that $\{f_{4,l}\}_{l \in A_{4,n}} \sim \{u_l\}_{l=1}^{|A_{4,n}|}$ for $n = 1, 2, \dots$; and by $F_4 = \overline{\text{span}}[f_{4,l}]_l$ denote the corresponding subspace.

To check conditions (7), it is enough to observe that for $i = 1, 2, 3$ and $m = 1, 2, \dots$, and every $K \in \mathcal{K}_{i+1,m}$ we have $K \subset A_{i+1,m}$; and hence $\{f_{i+1,l}\}_{l \in K} \sim \{u_l\}_{l=1}^{|K|}$, in other words, $F_{i+1}|_K \sim E|_K$. Therefore, by our construction,

$$\|I: F_{i+1}|_K \rightarrow F_i|_K\| \geq (1/2)\|I: E|_K \rightarrow F_i|_K\| \geq 2^{2m-1} = \delta_m^{-1}.$$

To check (5), note that for $i = 1, 2, 3$ and $m = 1, 2, \dots$ and every $A_{i,m}$ one has $F_i|_{A_{i,m}} \sim F_{i+1}|_{A_{i,m}} \sim E|_{A_{i,m}}$.

Finally, all the F_s 's are spanned by subsequences of the 1-unconditional basis, hence the projections $R_K^{(s)}$ have norms equal to 1. This shows case (Ia).

Case (Ib): Set $f_{1,l} = e_{1,l}$ for all l and let $\{A_{1,m}\}_m$ be singletons. Let $F_1 = \overline{\text{span}}[f_{1,l}]_l$. Since $\|I: E \rightarrow F_1\| \leq \|I : E \rightarrow X\| < \infty$, we must have $\|I : F_1 \rightarrow E\| = \infty$. Let $\Delta_2 = \{A_{2,m}\}_m$ be a partition of $\mathbb{N}$ into successive intervals such that $\|I: F_1|_{A_{2,m}} \rightarrow E|_{A_{2,m}}\| \geq 2^{2m}$ for all m . Define the sequence $\{f_{2,l}\}_l$ as a subsequence of $\{e_{2,l}\}_l$ in the same way as in Case (Ia) and let F_2 be its span. Then for all m we have $F_2|_{A_{2,m}} \sim E|_{A_{2,m}}$, hence $\|I: F_1|_{A_{2,m}} \rightarrow F_2|_{A_{2,m}}\| \geq 2^{2m}$.

Since $\|I: E \rightarrow F_1\| < \infty$, and $\|I: E \rightarrow F_2\| < \infty$, then also $\|I: E \rightarrow D(F_1 \oplus F_2)\| < \infty$, where $D(F_1 \oplus F_2) = \overline{\text{span}}[f_{1,l} + f_{2,l}]_l$. Thus $\|I: D(F_1 \oplus F_2) \rightarrow E\| = \infty$. As in Case (Ia) we can find an increasing sequence $\{M_j\}$ of integers and a partition $\Delta_3 = \{A_{3,j}\}_j$ of $\mathbb{N}$ with $A_{3,j} = \bigcup_{m=M_{j-1}+1}^{M_j} A_{2,m}$ such that for every $K \in \mathcal{K}_{3,j}$ we have $\|I: D(F_1 \oplus F_2)|_K \rightarrow E|_K\| \geq 2^{2j}$.

We then define a subsequence $\{f_{3,l}\}_l$ of $\{e_{3,l}\}_l$, so that its span F_3 satisfies $F_3|_{A_{3,j}} \sim E|_{A_{3,j}}$, for all j .

Finally we define a partition $\Delta_4 = \{A_{4,n}\}_n$ with $\Delta_4 \prec \Delta_3$ such that for every $K \in \mathcal{K}_{4,n}$ we have $\|I: D(F_1 \oplus F_2 \oplus F_3)|_K \rightarrow E|_K\| \geq 2^{2n}$ (for $n = 1, 2, \dots$), and then we define a subsequence $\{f_{4,l}\}_l$ of $\{e_{4,l}\}_l$, so that its span F_4 satisfies $F_4|_{A_{4,n}} \sim E|_{A_{4,n}}$, for all n .

Just as in Case (Ia), it is easy to check that (5) and (6) are satisfied, and that all the projections $R_K^{(s)}$ have norms equal to 1. $\square$

Proof of Theorem 3.1 in Case (II): The key argument in [K-T] was contained in Proposition 3.3. The same statement and the proof that followed, work here, provided that we can use Theorem $\star 2.1$. This requires that the notion of a C -regular pair $\{\Delta, F\}$ used in [K-T] should be replaced by a suitable modification, which we call here a C -projection-regular pair.

We say that a partition $\Delta = \{A_m\}_m$ of $\mathbb{N}$ into consecutive intervals and a space F with a normalized Schauder basis $\{f_l\}_l$ form a C -projection-regular pair, if the following conditions are satisfied:

- (i) $\text{equiv}(F|_{A_m}, l_2^{|A_m|}) \leq C$ and the natural projection $Q_{A_m}: F \rightarrow F|_{A_m}$ has the norm $\|Q_{A_m}\| \leq C$, for $m = 1, 2, \dots$;
- (ii) for every $L \in \mathcal{L}(\Delta)$, the basis $\{f_l\}_{l \in L}$ in $F|_L$ is 1-unconditional and the natural projection $Q_L: F \rightarrow F|_L$ has the norm $\|Q_L\| \leq C$ (here $\mathcal{L}(\Delta)$ is the family of all $L \subset \mathbb{N}$ such that $|L \cap A_m| = 1$ for all m).

(iii) for arbitrary $L, L' \in \mathcal{L}(\Delta)$ one has $\text{equiv}(F|_L, F|_{L'}) = 1$.

The modified Proposition 3.3 now additionally assumes that all spaces E_i have subsymmetric bases, and it asserts that the constructed pairs $\{\Delta_i, F_i\}$ are C -projection-regular.

The use of Rademacher functions in a setting of discrete Banach lattice gives more information for spaces with subsymmetric bases. For example, Tzafriri's argument mentioned in [K-T] uses this approach; this can be also tackled by a modification of inequalities from Lemma 3.2 in [K-T]. One gets that for $m \in \mathbb{N}$ and $N = 2^m$, if E is an N -dimensional space of cotype $r < \infty$ with a 1-subsymmetric basis $\{e_j\}_j$, then m Rademacher vectors $\{f_l\}$ of the form

$$f_l = \alpha \sum_{j=1}^N \varepsilon_l(j) e_j \quad (8)$$

are C -equivalent to the unit vector basis in ℓ_2^m , where C depends on r and the cotype r constant. Moreover, $\{\varepsilon_l(j)\}$ are values of Rademacher functions on the dyadic partition of the interval $[0, 1]$, and so they are mutually orthogonal, $\sum_j \varepsilon_l(j) \varepsilon_{l'}(j) = 0$ if $l \neq l'$.

For a given space E of cotype r with a subsymmetric basis $\{e_j\}_j$, and a partition $\Delta = \{A_m\}_m$ of $\mathbb{N}$ into consecutive intervals, we repeat word by word the first part of the proof of Proposition 3.3 in [K-T] to get a subspace $F = \overline{\text{span}}[f_l]_l$ of the following form: we fix arbitrary successive intervals of integers, $\{J_m\}$, with $|J_m| = 2^{|A_m|}$ and by formula (8) we define vectors f_l by

$$f_l = \alpha_m \sum_{j \in J_m} \varepsilon_l(j) e_j \quad \text{for } l \in A_m, \ m = 1, 2, \dots$$

All the conditions from [K-T] are satisfied as before, we need only to check the boundness of the natural projections, $Q_K: F \rightarrow F|_K$ given by $Q_K(\sum_l a_l f_l) = \sum_{l \in K} a_l f_l$, for $K \subset \mathbb{N}$.

For condition (i), Q_{A_m} is equal to the restriction to F of the natural projection P_{J_m} in E onto coordinates from J_m , hence is obviously bounded.

For condition (ii), let $L = \{l_m\}_m \in \mathcal{L}(\Delta)$. Then $\{f_{l_m}\}_m$ form a subspace of E spanned by constant coefficient successive blocks. Consider the averaging projection P_L in E given by the formula

$$P_L(\sum_j a_j e_j) = \sum_m (\alpha_m |J_m|)^{-1} \left(\sum_{j \in J_m} \varepsilon_{l_m}(j) a_j \right) f_{l_m}.$$

The orthogonality relations of sign vectors $\{\varepsilon_l(j)\}$ imply that for any $l \notin L$ we have $P_L(f_l) = 0$. So the projection Q_L on F is equal to the restriction of P_L to F . Finally let us recall (Lindenstrauss–Tzafriri, Vol I, 3.a.4) that averaging projections in a space with a subsymmetric basis have the norm $\|P_L\| \leq 2$, and this shows $\|Q_L\| \leq 2$.

This shows a modified version of Proposition 3.3 which allows to use Theorem $\star 2.1$ in the proof of Theorem 3.1. $\square$

Remark Actually, even in Case II, a subspace without l.u.st. can be constructed which has 2-dimensional unconditional decomposition. This can be done by a careful use of Krivine theorem on finite block representability of ℓ_p . Since the basis $\{e_l\}_l$ is subsymmetric, one can find $1 \leq p \leq \infty$ and an unconditional basic sequence $\{f_l\}_l$ in $X = \overline{\text{span}}[e_l]_l$ such that denoting by $\Delta = \{J_m\}_m$ the partition of $\mathbb{N}$ into successive intervals with $|J_m| = m$, we get $\{f_l\}_{l \in J_m} \sim \ell_p^m$ and for any $L, L' \in \mathcal{L}(\Delta)$, the sequences $\{f_l\}_{l \in L}$ and $\{f_l\}_{l \in L'}$ are 1-equivalent, but they are not equivalent to ℓ_p . In many cases, $\{f_l\}_l$ is a block basis consisting of Krivine's vectors; and then, for $l \in J_m$, the f_l 's are shifts of the same vector along the basis $\{e_l\}_l$. In remaining cases, for instance if $X = \ell_q$ for some $q \neq 2$, we let $p = 2$ and use Dvoretzky theorem. After constructing a suitable basis $\{f_l\}_l$, the construction follows lines similar as in Case I.

We feel, however, that the construction of Case II based on [K-T] may be of interest in other situations than Theorem 3.1 as well. Similarly, Theorem $\star 2.1$ is, as far as we know, one of very few results of this type which does not assume the unconditionality of the whole Schauder decomposition, but just of its subsets.

References:

[K-T] R. Komorowski & N. Tomczak-Jaegermann: Banach spaces without local unconditional structure; *Israel J. of Math.*, 89 (1995), pp. 205–226.

Institute of Mathematics, Technical University,
Wrocław, Poland,
`komo@graf.im.pwr.wroc.pl`

Department of Mathematics, University of Alberta,
Edmonton, Alberta, Canada T6G 2G1,
`ntomczak@approx.math.ualberta.ca`